


Patent claims

1. Compounds of the general formula (I)



in which

X represents O, S, SO, SO₂, CH₂, CHF, CF₂ or represents NR⁸ in which R⁸ represents hydrogen or (C₁-C₄)-alkyl,

R¹ and R² are identical or different and represent hydrogen or (C₁-C₄)-alkyl,

R³ and R⁴ are identical or different and represent hydrogen, halogen, cyano, (C₁-C₆)-alkyl, CF₃, CHF₂, CH₂F, vinyl or (C₃-C₇)-cycloalkyl, where at least one of the two substituents is not hydrogen,

R⁵ represents hydrogen, (C₁-C₄)-alkyl or halogen,

R⁶ represents a group of the formula -S-R⁹, -S(O)_n-R¹⁰, -NR¹¹-C(O)-R¹², -CH₂-R¹³ or -M-R¹⁴, in which

R⁹ represents (C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkyl, (C₂-C₆)-alkenyl, (C₆-C₁₀)-aryl, (C₆-C₁₀)-arylmethyl or represents a saturated, partially unsaturated or aromatic 5- to 10-membered heterocycle having up to four identical or different heteroatoms

from the group consisting of N, O and S, where the abovementioned radicals are optionally substituted by one, two or three identical or different substituents selected from the group consisting of halogen, hydroxyl, oxo, cyano, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, carboxyl and (C₁-C₄)-alkoxycarbonyl,

n represents the number 1 or 2,

R¹⁰ represents OR¹⁵, NR¹⁶R¹⁷, (C₁-C₁₀)-alkyl, (C₃-C₈)-cycloalkyl, (C₂-C₆)-alkenyl, (C₆-C₁₀)-aryl, (C₆-C₁₀)-arylmethyl or represents a saturated, partially unsaturated or aromatic 5- to 10-membered heterocycle having up to four identical or different heteroatoms from the group consisting of N, O and S, where the abovementioned radicals are optionally substituted by one, two or three identical or different substituents selected from the group consisting of halogen, hydroxyl, oxo, cyano, nitro, amino, NR¹⁸R¹⁹, trifluoromethyl, (C₁-C₆)-alkyl, optionally R²⁰-substituted (C₁-C₆)-alkoxy, (C₃-C₈)-cycloalkyl, (C₆-C₁₀)-aryl, which for its part is optionally substituted by halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, trifluoromethyl, nitro or cyano, -O-C(O)-R²¹, -C(O)-OR²², -C(O)-NR²³R²⁴, -SO₂-NR²⁵R²⁶, -NH-C(O)-R²⁷ and -NH-C(O)-OR²⁸, where

R¹⁵, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵, R²⁶, R²⁷ and R²⁸ are identical or different and each represents hydrogen, phenyl, benzyl, (C₁-C₆)-alkyl or (C₃-C₈)-cycloalkyl which for their part are optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, amino, carboxyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkoxycarbonyl, (C₁-

C₄)-alkoxycarbonylamino, (C₁-C₅)-alkanoyloxy, a heterocycle or by phenyl which for its part is optionally substituted by halogen or hydroxyl,

and

5

R¹⁶ and R¹⁷ are identical or different and independently of one another represent hydrogen, straight-chain or branched (C₁-C₆)-alkyl which may be mono- or polysubstituted by identical or different substituents from the group consisting of mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, (C₁-C₄)-alkoxy, (C₁-C₆)-alkoxycarbonyl, carboxyl, pyridyl or (C₆-C₁₀)-aryl, where the latter for its part is optionally substituted by halogen, tri-fluoromethyl, (C₁-C₆)-alkyl or (C₁-C₆)-alkoxy,

10

15

represent (C₃-C₈)-cycloalkyl or represent a 5- to 7-membered heterocycle which contains one or two nitrogen atoms, where cycloalkyl and heterocycle are optionally substituted by (C₁-C₄)-alkyl,

20

or

R¹⁶ and R¹⁷ together with the nitrogen atom to which they are attached form a 5- to 7-membered saturated, optionally benzo-fused heterocycle which may contain up to two further heteroatoms from the group consisting of N, O and S and which may be substituted by amino, (C₁-C₆)-alkyl, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino or phenyl,

25

30

R¹¹ represents hydrogen, straight-chain or branched (C₁-C₆)-alkyl which may be mono- or polysubstituted by identical or different substituents from the group consisting of mono-(C₁-C₆)-alkylamino, di-(C₁-C₆)-alkylamino, (C₁-C₄)-alkoxy, (C₁-C₆)-alkoxycarbonyl, carboxyl, pyridyl and (C₆-C₁₀)-aryl, where the latter for its part is optionally substituted by halogen, trifluoromethyl, (C₁-C₆)-alkyl or (C₁-C₆)-alkoxy, represents (C₃-C₈)-cycloalkyl or represents a 5- to 7-membered heterocycle which contains one or two nitrogen atoms, where cycloalkyl and heterocycle are optionally substituted by (C₁-C₄)-alkyl,

R¹² represents straight-chain or branched (C₁-C₁₅)-alkyl which may be substituted by (C₃-C₈)-cycloalkyl, (C₁-C₄)-alkoxy, phenyl, phenoxy or benzyloxy, where the aromatic radicals mentioned for their part may each be substituted up to three times by identical or different substituents from the group consisting of halogen, (C₁-C₆)-alkyl and (C₁-C₄)-alkoxy,

represents (C₃-C₈)-cycloalkyl which may be substituted by (C₁-C₄)-alkoxy or phenyl,

represents (C₆-C₁₀)-aryl which may be substituted up to three times by identical or different substituents from the group consisting of (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy, halogen, cyano, amino, trifluoromethyl and phenyl,

or

represents a 5- or 6-membered saturated or aromatic, optionally benzo-fused heterocycle having up to two heteroatoms from the group consisting of N, O and S,

5

or

represents a group of the formula $-OR^{29}$ or $-NR^{30}R^{31}$,

in which

10

R^{29} represents straight-chain or branched (C_1-C_6) -alkyl,

and

15

R^{30} and R^{31} are identical or different and independently of one another

represent hydrogen, straight-chain or branched (C_1-C_{12}) -alkyl which may be substituted by aminocarbonyl, a group of the formula $-NR^{32}R^{33}$, 5- or 6-membered heteroaryl which contains up to 3 heteroatoms selected from the group consisting of N, O and S, or by phenyl, where phenyl is optionally substituted up to two times by identical or different substituents from the group consisting of halogen, (C_1-C_4) -alkyl, trifluoromethyl and (C_1-C_4) -alkoxy,

20

25

represent (C_3-C_8) -cycloalkyl which may be substituted by (C_1-C_4) -alkyl,

30

- 77 -

represent (C₆-C₁₀)-aryl which may be substituted up to three times by identical or different substituents from the group consisting of halogen, (C₁-C₄)-alkyl, trifluoromethyl, (C₁-C₄)-alkoxy, amino, phenyl and phenoxy,

or

represent a 5- to 7-membered saturated or unsaturated heterocycle which contains one or two nitrogen atoms and is optionally substituted by (C₁-C₄)-alkyl or an oxo group,

where

R³² and R³³ are identical or different and independently of one another represent hydrogen, (C₁-C₆)-alkyl, phenyl or (C₆-C₁₀)-arylsulphonyl,

or

together with the nitrogen atom to which they are attached form a 3- to 7-membered saturated heterocycle which optionally contains up to two further heteroatoms from the group consisting of N, O and S,

or

R³⁰ and R³¹ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle which may contain up to two further heteroatoms from

the group consisting of N, O and S and which may be substituted by amino, (C₁-C₆)-alkyl, (C₁-C₄)-alkanoyl, aminocarbonyl, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino, phenyl or pyridyl,

5

R¹³ represents a saturated, partially unsaturated or aromatic 5- to 10-membered heterocycle having up to three identical or different heteroatoms from the group consisting of N, O and S, which is optionally substituted by one, two or three identical or different substituents selected from the group consisting of (C₁-C₄)-alkyl, hydroxyl, oxo, (C₁-C₄)-alkoxy, halogen, cyano, carboxyl and (C₁-C₄)-alkoxycarbonyl, with the proviso that X in this case does not represent SO or SO₂,

10

15

or

R¹³ represents the group -NR³⁴R³⁵ in which

20

R³⁴ and R³⁵ are identical or different and represent hydrogen, (C₁-C₈)-alkyl which may be substituted by (C₆-C₁₀)-aryl, represent (C₃-C₈)-cycloalkyl, (C₆-C₁₀)-aryl or represent 5- or 6-membered heteroaryl having up to three identical or different heteroatoms from the group consisting of N, O and S where aryl and heteroaryl for their part are in each case optionally mono- or disubstituted by identical or different substituents from the group consisting of hydroxyl, amino, cyano, halogen, trifluoromethyl, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, carboxyl, (C₁-C₄)-alkoxycarbonyl and mono- and di-(C₁-C₄)-alkylaminocarbonyl,

25

30

M represents C=O, CH(OH), CHF or CF₂,

and

5


R¹⁴ has the meaning of R¹⁰ given above,

R⁷ represents hydrogen, (C₁-C₄)-alkyl or (C₁-C₄)-alkanoyl,

10

and

Z represents a group of the formula



in which

15

A represents O or S,

a represents the number 0 or 1,

20

D represents a straight-chain (C₁-C₄)-alkylene group which may be mono- or polysubstituted by identical or different substituents from the group consisting of (C₁-C₃)-alkyl, hydroxyl and fluorine,

25

and

R³⁶ represents OR³⁷ or NR³⁸R³⁹, in which

R^{37} , R^{38} and R^{39} are identical or different and each represents hydrogen, phenyl, benzyl, (C₁-C₆)-alkyl or (C₃-C₈)-cycloalkyl which for their part are optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, amino, carboxyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino, (C₁-C₅)-alkanoyloxy, a heterocycle or by phenyl which for its part is optionally substituted by halogen or hydroxyl,

and their pharmaceutically acceptable salts, solvates, hydrates and hydrates of the salts.

2. Compounds of the general formula (I) according to Claim 1

in which

X represents O, S, CH₂ or CF₂,

R^1 and R^2 are identical or different and represent hydrogen or methyl,

R^3 and R^4 are identical or different and represent hydrogen, halogen, (C₁-C₄)-alkyl, CF₃, CHF₂, CH₂F, vinyl or (C₃-C₅)-cycloalkyl, where at least one of the two substituents is not hydrogen,

R^5 represents hydrogen, (C₁-C₃)-alkyl, fluorine, chlorine or bromine,

R^6 represents a group of the formula -S(O)₂-R¹⁰, -NR¹¹-C(O)-R¹², -CH₂-R¹³ or -M-R¹⁴, in which

5 R^{10} represents $NR^{16}R^{17}$, (C_1-C_8) -alkyl, (C_5-C_7) -cycloalkyl, phenyl, benzyl or represents a saturated, partially unsaturated or aromatic 5- to 10-membered heterocycle having up to three identical or different heteroatoms from the group consisting of N, O and S, where the abovementioned radicals are optionally substituted by one, two or three identical or different substituents selected from the group consisting of halogen, hydroxyl, oxo, cyano, nitro, amino, dimethylamino, trifluoromethyl, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, (C_3-C_6) -cycloalkyl, phenyl, which for its parts is optionally substituted by halogen, (C_1-C_4) -alkyl, (C_1-C_4) -alkoxy, trifluoromethyl, nitro or cyano, $-C(O)-OR^{22}$, $-C(O)-NR^{23}R^{24}$, $-SO_2-NR^{25}R^{26}$, $-NH-C(O)-R^{27}$ and $-NH-C(O)-OR^{28}$, where

15 R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} and R^{28} are identical or different and each represents hydrogen, phenyl, benzyl, (C_1-C_4) -alkyl or (C_5-C_7) -cycloalkyl which for their part are optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, amino, carboxyl, (C_1-C_4) -alkoxy, (C_1-C_4) -alkoxycarbonyl, (C_1-C_4) -alkoxycarbonylamino or (C_1-C_5) -alkanoyloxy,

20 and

25 R^{16} and R^{17} are identical or different and independently of one another represent hydrogen, straight-chain or branched (C_1-C_6) -alkyl which may be mono- or polysubstituted by identical or different substituents from the group consisting of (C_1-C_4) -alkoxy, (C_1-C_4) -alkoxycarbonyl,

30

carboxyl, pyridyl or phenyl, where the latter for its part is optionally substituted by halogen, trifluoromethyl, (C₁-C₄)-alkyl or (C₁-C₄)-alkoxy,

5 represent (C₅-C₇)-cycloalkyl or represent a 5- to 7-membered heterocycle which contains one or two nitrogen atoms, where cycloalkyl and heterocycle are optionally substituted by (C₁-C₄)-alkyl,

10 or

R¹⁶ and R¹⁷ together with the nitrogen atom to which they are attached form a 5- to 7-membered saturated heterocycle which may contain up to two further heteroatoms from the group consisting of N, O and S and which may be substituted by amino, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino or phenyl,

20 R¹¹ represents hydrogen, straight-chain or branched (C₁-C₄)-alkyl, benzyl, (C₃-C₇)-cycloalkyl or represents a 5- to 7-membered heterocycle which contains one or two nitrogen atoms, where cycloalkyl and heterocycle are optionally substituted by (C₁-C₄)-alkyl,

25 R¹² represents straight-chain or branched (C₁-C₈)-alkyl which may be substituted by (C₃-C₇)-cycloalkyl, (C₁-C₄)-alkoxy, phenyl, phenoxy or benzyloxy, where the aromatic radicals mentioned for their part may each be substituted up to three times by

identical or different substituents from the group consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkoxy,

or

5

represents phenyl which may be substituted up to three times by identical or different substituents from the group consisting of (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, halogen, cyano, amino and trifluoromethyl,

10

or

represents a group of the formula -OR²⁹ or -NR³⁰R³¹,
in which

15

R²⁹ represents straight-chain or branched (C₁-C₄)-alkyl,

and

20

R³⁰ and R³¹ are identical or different and independently of one another

25

represent hydrogen, straight-chain or branched (C₁-C₈)-alkyl which may be substituted by phenyl, which for its part is optionally substituted up to two times by identical or different substituents from the group consisting of halogen, (C₁-C₄)-alkyl, trifluoromethyl and (C₁-C₄)-alkoxy,

- 84 -

represent (C₃-C₇)-cycloalkyl which may be substituted by (C₁-C₄)-alkyl,

or

represent phenyl which may be substituted up to three times by identical or different substituents from the group consisting of halogen, (C₁-C₄)-alkyl, trifluoromethyl, (C₁-C₄)-alkoxy and amino,

or

R³⁰ and R³¹ together with the nitrogen atom to which they are attached form a 5- to 7-membered saturated heterocycle which may contain up to two further heteroatoms from the group consisting of N, O and S and which may be substituted by amino, (C₁-C₄)-alkyl, (C₁-C₄)-alkanoyl, aminocarbonyl, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino or phenyl,

R¹³ represents a saturated, partially unsaturated or aromatic 5- or 6-membered heterocycle having up to three identical or different heteroatoms from the group consisting of N, O and S, which is optionally substituted by one, two or three identical or different substituents selected from the group consisting of (C₁-C₄)-alkyl, hydroxyl, oxo, (C₁-C₄)-alkoxy, halogen, cyano, carboxyl and (C₁-C₄)-alkoxycarbonyl,

or

represents the group $-\text{NR}^{34}\text{R}^{35}$, in which

R^{34} and R^{35} are identical or different and represent hydrogen, $(\text{C}_1\text{-C}_6)$ -alkyl, which may be substituted by phenyl, represent $(\text{C}_5\text{-C}_7)$ -cycloalkyl, phenyl or represent 5- or 6-membered heteroaryl having up to three identical or different heteroatoms from the group consisting of N, O and S, where phenyl and heteroaryl for their part are each optionally mono- or disubstituted by identical or different substituents from the group consisting of hydroxyl, amino, cyano, halogen, $(\text{C}_1\text{-C}_4)$ -alkyl, trifluoromethyl, $(\text{C}_1\text{-C}_4)$ -alkoxy, carboxyl or $(\text{C}_1\text{-C}_4)$ -alkoxycarbonyl,

M represents $\text{C}=\text{O}$, $\text{CH}(\text{OH})$ or CF_2 ,


and

R^{14} has the meaning of R^{10} given above,

R^7 represents hydrogen, methyl or acetyl,

and

Z represents a group of the formula



in which

A represents O or S,

a represents the number 0 or 1,

5 D represents a straight-chain (C₁-C₃)-alkylene group which may be mono- or polysubstituted by identical or different substituents from the group consisting of methyl, hydroxyl and fluorine,

and

10

R³⁶ represents OR³⁷ or NR³⁸R³⁹, in which

15

R³⁷ represents hydrogen, phenyl, benzyl, (C₁-C₆)-alkyl or (C₃-C₇)-cycloalkyl which for their part are optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, amino, carboxyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino, (C₁-C₅)-alkanoyloxy and a heterocycle,

20

and

25

R³⁸ and R³⁹ are identical or different and each represents hydrogen, (C₁-C₆)-alkyl or (C₃-C₇)-cycloalkyl, which for their part are optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, amino, carboxyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkoxycarbonylamino, (C₁-C₅)-alkanoyloxy, a heterocycle and phenyl which for its part is optionally substituted by halogen or hydroxyl,

30

and their pharmaceutically acceptable salts, solvates, hydrates and hydrates of the salts.

3. Compounds of the general formula (I) according to Claim 1

5

in which

X represents O, S or CH₂,

10

R¹ and R² represent hydrogen,

R³ and R⁴ are identical or different and represent methyl, ethyl, propyl, isopropyl, cyclopropyl, trifluoromethyl, chlorine or bromine,

15

R⁵ represents hydrogen,

R⁶ represents a group of the formula -S(O)₂-R¹⁰, -NH-C(O)-R¹², -CH₂-R¹³, -C(O)-R¹⁴ or -CH(OH)-R⁴⁰, in which

20

R¹⁰ represents phenyl or represents 5- or 6-membered heteroaryl having up to three identical or different heteroatoms from the group consisting of N, O and S, which radicals are optionally mono- or disubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine, hydroxyl, cyano, trifluoromethyl, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, carboxyl and (C₁-C₄)-alkoxycarbonyl,

25

or

30

represents the group -NR¹⁶R¹⁷, in which

R^{16} and R^{17} together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocycle which may contain a further heteroatom from the group consisting of N, O and S and which may be substituted by (C_1-C_4) -alkyl,

R^{12} represents straight-chain or branched (C_1-C_6) -alkyl which is optionally substituted by phenoxy or benzyloxy,

R^{13} represents 5- or 6-membered heteroaryl having up to three identical or different heteroatoms from the group consisting of N, O and S, which is optionally substituted by one or two identical or different substituents selected from the group consisting of (C_1-C_4) -alkyl, hydroxyl, (C_1-C_4) -alkoxy, fluorine, chlorine, bromine, cyano, carboxyl and (C_1-C_4) -alkoxycarbonyl, or represents the group $-NR^{34}R^{35}$, in which R^{34} represents (C_1-C_6) -alkyl or (C_5-C_7) -cycloalkyl,

and

R^{35} represents benzyl which is optionally substituted in the phenyl ring by hydroxyl, (C_1-C_4) -alkoxy, (C_1-C_4) -alkyl, trifluoromethyl, fluorine, chlorine or cyano,

R^{14} represents a group of the formula $-NR^{41}R^{42}$, in which

R^{41} represents hydrogen, (C_1-C_6) -alkyl or (C_3-C_7) -cycloalkyl,

R^{42} represents hydrogen or represents (C₁-C₄)-alkyl which may be substituted by phenyl,

or

R^{41} and R^{42} together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocycle which may contain a further heteroatom from the group consisting of N, O and S and which may be substituted by (C₁-C₄)-alkyl,

and


R^{40} represents phenyl or naphthyl, which are optionally mono- or disubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, cyano, trifluoromethyl or (C₁-C₄)-alkoxycarbonyl,

R^7 represents hydrogen,

and

Z represents a group of the formula

- 90 -



in which R^{36} represents hydroxyl or the radical $-\text{C}(\text{O})-\text{R}^{36}$ has the meanings of R^{36} given above for a group which, in the sense of a prodrug, can be degraded to the carboxylic acid $-\text{C}(\text{O})-\text{OH}$ or a salt thereof,

and their pharmaceutically acceptable salts, solvates, hydrates and hydrates of the salts.

4. Compounds of the general formula (I) according to Claim 1

in which

X represents CH_2 or, in particular, oxygen,

R^1 and R^2 represent hydrogen,

R^3 and R^4 are identical or different and represent methyl, ethyl, propyl, isopropyl, cyclopropyl, trifluoromethyl, chlorine or bromine,


R^5 represents hydrogen,

R^6 represents a group of the formula $-S(O)_2-R^{10}$, $-CH_2-R^{13}$ or $-C(O)-R^{14}$,
in which

R^{10} represents phenyl, pyridyl, pyrimidinyl or pyridazinyl which
are optionally mono- or disubstituted by identical or different
substituents from the group consisting of fluorine, chlorine,
bromine, hydroxyl, cyano, trifluoromethyl, (C_1-C_4) -alkyl, $(C_1-$
 $C_4)$ -alkoxy, carboxyl and (C_1-C_4) -alkoxycarbonyl,

or

represents a group of the formula



R^{13} represents pyridyl, pyrimidinyl or pyridazinyl which are
optionally substituted by one or two identical or different
substituents selected from the group consisting of (C_1-C_4) -
alkyl, hydroxyl, (C_1-C_4) -alkoxy, fluorine, chlorine, bromine,
cyano, carboxyl and (C_1-C_4) -alkoxycarbonyl, or represents the
group $-NR^{34}R^{35}$, in which

R^{34} represents (C_1-C_4) -alkyl or (C_5-C_7) -cycloalkyl,

and

R^{35} represents benzyl which is optionally substituted in the phenyl ring by hydroxyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkyl, trifluoromethyl, fluorine, chlorine or cyano,

5 and

R^{14} represents a group of the formula $-NR^{41}R^{42}$, in which

R^{41} represents hydrogen, (C₁-C₄)-alkyl or (C₅-C₇)-cycloalkyl,

and


R^{42} represents hydrogen or represents (C₁-C₄)-alkyl which may be substituted by phenyl,

R^7 represents hydrogen,

and

Z represents a group of the formula


- 93 -



in which R^{37} represents hydrogen, $(\text{C}_1\text{-C}_4)$ -alkyl or $(\text{C}_4\text{-C}_6)$ -cycloalkyl,

and their pharmaceutically acceptable salts, solvates, hydrates and hydrates of the salts.

5. Compounds of the formula (Ia)



in which

X represents CH_2 or O,

R^3 and R^4 are identical or different and represent bromine, trifluoromethyl, ethyl, cyclopropyl and, in particular, methyl or chlorine,


Z represents a group of the formula $-\text{CH}_2-\text{C}(\text{O})-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{C}(\text{O})-\text{OH}$, $-\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{OH}$ or $-\text{S}-\text{CH}_2-\text{C}(\text{O})-\text{OH}$,

and

R^6 represents a group of the formula $-S(O)_2-R^{10}$, in which

R^{10} represents phenyl or represents pyridyl which are optionally mono- or disubstituted by identical or different substituents from the group consisting of fluorine, chlorine, cyano, trifluoromethyl, methyl, hydroxyl and methoxy.

6. Compounds of the formula (Ia)



in which

X represents CH_2 or O,

R^3 and R^4 are identical or different and represent bromine, trifluoromethyl, ethyl, cyclopropyl and, in particular, methyl or chlorine,

Z represents a group of the formula $-CH_2-C(O)-OH$, $-CH_2-CH_2-C(O)-OH$, $-O-CH_2-C(O)-OH$, $-O-C[(CH_3)_2]-C(O)-OH$ or $-S-CH_2-C(O)-OH$,

and

R^6 represents a group of the formula $-S(O)_2-R^{10}$, in which

R¹⁰ represents phenyl or represents pyridyl which are optionally mono- or disubstituted by identical or different substituents from the group consisting of fluorine, chlorine, cyano, trifluoromethyl, methyl, hydroxyl and methoxy.

5

7. Compounds as defined in Claims 1 to 6 for preventing and treating diseases.

8. Medicaments, comprising at least one compound of the general formula (I) as defined in Claims 1 to 6.

10

9. Process for preparing medicaments, characterized in that at least one compound of the general formula (I) as defined in Claims 1 to 6 is converted, using excipients and/or carriers, into a suitable administration form.

15

10. Use of compounds of the general formula (I) as defined in Claims 1 to 6 for preparing medicaments.

11. Use according to Claim 8 for preparing medicaments for the treatment and/or prophylaxis of arteriosclerosis, obesity and/or hypercholesterolaemia.

20

12. Use according to Claim 8 for preparing medicaments for the prophylaxis and/or treatment of disease forms which can be treated with natural thyroid hormone.


25

13. Method for the treatment and/or prophylaxis of disorders, characterized in that compounds as defined in Claims 1 to 6 are used.

14. Process for preparing compounds of the formula (I) as defined in Claim 1, characterized in that reactive phenol derivatives of the general formula (II)

30

- 96 -



in which


5 R⁵ and R⁶ are as defined in Claim 1 and

PG represents a protective group and

V represents a binding or leaving group,

10

are reacted, if appropriate with isolation of the intermediates, or directly, with reactive phenyl derivatives of the general formula (III)



15

in which

R¹, R², R³ and R⁴ are as defined in Claim 1 and

W represents a binding or leaving group and

20

Z' has the meaning given for Z or represents OH, O-PG, SH, S-PG, or represents an aldehyde, cyano, carboxyl or (C₁-C₄)-alkoxycarbonyl group,

if appropriate in the presence of inert solvents and catalysts, to give compounds of the formula (I).

姓名	性别	年龄	职业	住址	联系电话	备注
王德胜	男	45	教师	XX市XX区XX路XX号	138XXXXXXX	
李小红	女	32	护士	XX市XX区XX路XX号	139XXXXXXX	
张小明	男	28	程序员	XX市XX区XX路XX号	137XXXXXXX	
赵大伟	男	50	工程师	XX市XX区XX路XX号	136XXXXXXX	
陈丽娟	女	40	医生	XX市XX区XX路XX号	135XXXXXXX	
刘国强	男	35	经理	XX市XX区XX路XX号	134XXXXXXX	
孙秀英	女	55	退休	XX市XX区XX路XX号	133XXXXXXX	
周志远	男	42	律师	XX市XX区XX路XX号	132XXXXXXX	
吴小芳	女	30	会计	XX市XX区XX路XX号	131XXXXXXX	
郑大刚	男	48	司机	XX市XX区XX路XX号	130XXXXXXX	
冯娟娟	女	38	销售	XX市XX区XX路XX号	129XXXXXXX	
马志强	男	52	教授	XX市XX区XX路XX号	128XXXXXXX	
徐小华	女	25	学生	XX市XX区XX路XX号	127XXXXXXX	
黄大伟	男	40	工人	XX市XX区XX路XX号	126XXXXXXX	
曹丽娟	女	35	文员	XX市XX区XX路XX号	125XXXXXXX	
周志远	男	50	老板	XX市XX区XX路XX号	124XXXXXXX	
吴小芳	女	30	护士	XX市XX区XX路XX号	123XXXXXXX	
郑大刚	男	45	教师	XX市XX区XX路XX号	122XXXXXXX	
冯娟娟	女	38	医生	XX市XX区XX路XX号	121XXXXXXX	
马志强	男	52	工程师	XX市XX区XX路XX号	120XXXXXXX	
徐小华	女	25	程序员	XX市XX区XX路XX号	119XXXXXXX	
黄大伟	男	40	经理	XX市XX区XX路XX号	118XXXXXXX	
曹丽娟	女	35	会计	XX市XX区XX路XX号	117XXXXXXX	
周志远	男	50	司机	XX市XX区XX路XX号	116XXXXXXX	
吴小芳	女	30	销售	XX市XX区XX路XX号	115XXXXXXX	
郑大刚	男	45	教授	XX市XX区XX路XX号	114XXXXXXX	
冯娟娟	女	38	学生	XX市XX区XX路XX号	113XXXXXXX	
马志强	男	52	工人	XX市XX区XX路XX号	112XXXXXXX	
徐小华	女	25	文员	XX市XX区XX路XX号	111XXXXXXX	
黄大伟	男	40	老板	XX市XX区XX路XX号	110XXXXXXX	
曹丽娟	女	35	护士	XX市XX区XX路XX号	109XXXXXXX	
周志远	男	50	教师	XX市XX区XX路XX号	108XXXXXXX	
吴小芳	女	30	医生	XX市XX区XX路XX号	107XXXXXXX	
郑大刚	男	45	工程师	XX市XX区XX路XX号	106XXXXXXX	
冯娟娟	女	38	程序员	XX市XX区XX路XX号	105XXXXXXX	
马志强	男	52	经理	XX市XX区XX路XX号	104XXXXXXX	
徐小华	女	25	会计	XX市XX区XX路XX号	103XXXXXXX	
黄大伟	男	40	司机	XX市XX区XX路XX号	102XXXXXXX	
曹丽娟	女	35	销售	XX市XX区XX路XX号	101XXXXXXX	
周志远	男	50	教授	XX市XX区XX路XX号	100XXXXXXX	
吴小芳	女	30	学生	XX市XX区XX路XX号	99XXXXXXX	
郑大刚	男	45	工人	XX市XX区XX路XX号	98XXXXXXX	
冯娟娟	女	38	文员	XX市XX区XX路XX号	97XXXXXXX	
马志强	男	52	老板	XX市XX区XX路XX号	96XXXXXXX	
徐小华	女	25	护士	XX市XX区XX路XX号	95XXXXXXX	
黄大伟	男	40	教师	XX市XX区XX路XX号	94XXXXXXX	
曹丽娟	女	35	医生	XX市XX区XX路XX号	93XXXXXXX	
周志远	男	50	工程师	XX市XX区XX路XX号	92XXXXXXX	
吴小芳	女	30	程序员	XX市XX区XX路XX号	91XXXXXXX	
郑大刚	男	45	经理	XX市XX区XX路XX号	90XXXXXXX	
冯娟娟	女	38	会计	XX市XX区XX路XX号	89XXXXXXX	
马志强	男	52	司机	XX市XX区XX路XX号	88XXXXXXX	
徐小华	女	25	销售	XX市XX区XX路XX号	87XXXXXXX	
黄大伟	男	40	教授	XX市XX区XX路XX号	86XXXXXXX	
曹丽娟	女	35	学生	XX市XX区XX路XX号	85XXXXXXX	
周志远	男	50	工人	XX市XX区XX路XX号	84XXXXXXX	
吴小芳	女	30	文员	XX市XX区XX路XX号	83XXXXXXX	
郑大刚	男	45	老板	XX市XX区XX路XX号	82XXXXXXX	
冯娟娟	女	38	护士	XX市XX区XX路XX号	81XXXXXXX	
马志强	男	52	教师	XX市XX区XX		